



Knowledge grows

Late winterlick for cattle and sheep

Late dry season

These licks are suitable for usage in the late dry season when the quality of grazing is at its lowest. This usually coincides with an overall shortage in available grazing. These two factors lead to low dry-matter intakes that prohibit production. For optimal reproduction it is essential to maintain the body condition of reproducing animals during this period. The correct supplementation of protein, energy and minerals is required.

Protein supplementation for late dry season for cattle and sheep

| Item | kg/ mixture |
|----------------------------------|------------------|
| Maize meal | 300 |
| Oilcake | 300 |
| Feed Grade Urea | 100 |
| Kimtrafos 12 Grandé/PhosSure 12 | 75 |
| Kalori 3000 | 25 |
| Feed Grade Sulphur | 5 |
| Salt | 300 |
| TOTAL | 1105 |
| Composition | |
| Crude Protein(%) | 38.7 |
| ME (MJ/ kg) | 6.6 |
| Ca (%) | 1.8 |
| P (%) | 1.1 |
| S (%) | 0.5 |
| INTAKE, g/ day cattle | 500 – 600 |
| INTAKE, g/sheep, goat/day | 150 - 170 |

Supplemented under the following circumstances:

- Particularly suitable for mid-/late-pregnant beef cows without calves in a poor condition.

Tel: +27 31 910 5100

Mail: animal.nutrition.sa@yara.com

